

Every function is the representation function of an additive basis for the integers*

Melvyn B. Nathanson[†]
 Department of Mathematics
 Lehman College (CUNY)
 Bronx, New York 10468
 Email: nathansn@alpha.lehman.cuny.edu

Abstract

Let A be a set of integers. For every integer n , let $r_{A,h}(n)$ denote the number of representations of n in the form $n = a_1 + a_2 + \cdots + a_h$, where $a_1, a_2, \dots, a_h \in A$ and $a_1 \leq a_2 \leq \cdots \leq a_h$. The function

$$r_{A,h} : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$$

is the *representation function of order h for A* . The set A is called an *asymptotic basis of order h* if $r_{A,h}^{-1}(0)$ is finite, that is, if every integer with at most a finite number of exceptions can be represented as the sum of exactly h not necessarily distinct elements of A . It is proved that every function is a representation function, that is, if $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ is any function such that $f^{-1}(0)$ is finite, then there exists a set A of integers such that $f(n) = r_{A,h}(n)$ for all $n \in \mathbf{Z}$. Moreover, the set A can be arbitrarily sparse in the sense that, if $\varphi(x) \geq 0$ for $x \geq 0$ and $\varphi(x) \rightarrow \infty$, then there exists a set A with $f(n) = r_{A,h}(n)$ and $\text{card}(\{a \in A : |a| \leq x\}) < \varphi(x)$ for all x .

It is an open problem to construct dense sets of integers with a prescribed representation function. Other open problems are also discussed.

1 Additive bases and the Erdős-Turán conjecture

Let $\mathbf{N}$, $\mathbf{N}_0$, and $\mathbf{Z}$ denote the positive integers, nonnegative integers, and integers, respectively. Let A be a set of integers. For every positive integer h , we define the *sumset*

$$hA = \{a_1 + \cdots + a_h : a_i \in A \text{ for all } i = 1, \dots, h\}.$$

*2000 Mathematics Subject Classification: 11B13, 11B34, 11B05. Key words and phrases. Additive bases, sumsets, representation functions, density, Erdős-Turán conjecture, Sidon set.

[†]This work was supported in part by grants from the NSA Mathematical Sciences Program and the PSC-CUNY Research Award Program.

We denote by $r_{A,h}(n)$ the number of representations of n in the form $n = a_1 + a_2 + \cdots + a_h$, where $a_1, a_2, \dots, a_h \in A$ and $a_1 \leq a_2 \leq \cdots \leq a_h$. The function $r_{A,h}$ is called the *representation function of order h* of the set A .

We denote the cardinality of the set A by $\text{card}(A)$. If A is a finite set of integers, we denote the maximum element of A by $\max(A)$. For any integer t and set $A \subseteq \mathbf{Z}$, we define the *translate*

$$t + A = \{t + a : a \in A\}.$$

In this paper we consider additive bases for the set of all integers. The set A of integers is called a *basis of order h for $\mathbf{Z}$* if every integer can be represented as the sum of h not necessarily distinct elements of A . The set A of integers is called an *asymptotic basis of order h for $\mathbf{Z}$* if every integer with at most a finite number of exceptions can be represented as the sum of h not necessarily distinct elements of A . Equivalently, the set A is an asymptotic basis of order h if the representation function $r_{A,h} : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ satisfies the condition

$$\text{card} \left(r_{A,h}^{-1}(0) \right) < \infty.$$

For any set X , let $\mathcal{F}_0(X)$ denote the set of all functions

$$f : X \rightarrow \mathbf{N}_0 \cup \{\infty\}$$

such that

$$\text{card} \left(f^{-1}(0) \right) < \infty.$$

We ask: Which functions in $\mathcal{F}_0(\mathbf{Z})$ are representation functions of asymptotic bases for the integers? This question has a remarkably simple and surprising answer. In the case $h = 1$ we observe that $f \in \mathcal{F}_0(\mathbf{Z})$ is a representation function if and only if $f(n) = 1$ for all integers $n \notin f^{-1}(0)$. For $h \geq 2$ we shall prove that *every* function in $\mathcal{F}_0(\mathbf{Z})$ is a representation function. Indeed, if $f \in \mathcal{F}_0(\mathbf{Z})$ and $h \geq 2$, then there exist infinitely many sets A such that $f(n) = r_{A,h}(n)$ for every $n \in \mathbf{Z}$. Moreover, we shall prove that we can construct arbitrarily sparse asymptotic bases A with this property. Nathanson [7] previously proved this theorem for $h = 2$ and the function $f(n) = 1$ for all $n \in \mathbf{Z}$.

This result about asymptotic bases for the integers contrasts sharply with the case of the nonnegative integers. The set A of nonnegative integers is called an *asymptotic basis of order h for $\mathbf{N}_0$* if every sufficiently large integer can be represented as the sum of h not necessarily distinct elements of A . Very little is known about the class of representation functions of asymptotic bases for $\mathbf{N}_0$. However, if $f \in \mathcal{F}_0(\mathbf{N}_0)$, then Nathanson [5] proved that there exists at most one set A such that $r_{A,h}(n) = f(n)$.

Many of the results that have been proved about asymptotic bases for $\mathbf{N}_0$ are negative. For example, Dirac [2] showed that the representation function of an asymptotic basis of order 2 cannot be eventually constant. Erdős and Fuchs [4] proved that the average value of a representation function of order 2

cannot even be approximately constant, in the sense that, for every infinite set A of nonnegative integers and every real number $c > 0$,

$$\sum_{n \leq N} r_{A,2}(n) \neq cN + o\left(N^{1/4} \log^{-1/2} N\right).$$

Erdős and Turán [3] conjectured that if A is an asymptotic basis of order h for the nonnegative integers, then the representation function $r_{A,h}(n)$ must be unbounded, that is,

$$\limsup_{n \rightarrow \infty} r_{A,h}(n) = \infty.$$

This famous unsolved problem in additive number theory is only a special case of the general problem of classifying the representation functions of asymptotic bases of finite order for the nonnegative integers.

2 Two lemmas

We use the following notation. For sets A and B of integers and for any integer t , we define the *sumset*

$$A + B = \{a + b : a \in A, b \in B\},$$

the *translation*

$$A + t = \{a + t : a \in A\},$$

and the *difference set*

$$A - B = \{a - b : a \in A, b \in B\}.$$

For every nonnegative integer h we define the *h -fold sumset* hA by induction:

$$0A = \{0\},$$

$$hA = A + (h-1)A = \{a_1 + a_2 + \cdots + a_h : a_1, a_2, \dots, a_h \in A\}.$$

The *counting function* for the set A is

$$A(y, x) = \text{card}(\{a \in A : y \leq a \leq x\}).$$

In particular, $A(-x, x)$ counts the number of integers $a \in A$ with $|a| \leq x$.

Let $[x]$ denote the integer part of the real number x .

Lemma 1 *Let $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ be a function such that $f^{-1}(0)$ is finite. Let Δ denote the cardinality of the set $f^{-1}(0)$. Then there exists a sequence $U = \{u_k\}_{k=1}^{\infty}$ of integers such that, for every $n \in \mathbf{Z}$ and $k \in \mathbf{N}$,*

$$f(n) = \text{card}(\{k \geq 1 : u_k = n\})$$

and

$$|u_k| \leq \left\lceil \frac{k + \Delta}{2} \right\rceil.$$

Proof. Every positive integer m can be written uniquely in the form

$$m = s^2 + s + 1 + r,$$

where s is a nonnegative integer and $|r| \leq s$. We construct the sequence

$$\begin{aligned} V &= \{0, -1, 0, 1, -2, -1, 0, 1, 2, -3, -2, -1, 0, 1, 2, 3, \dots\} \\ &= \{v_m\}_{m=1}^{\infty}, \end{aligned}$$

where

$$v_{s^2+s+1+r} = r \quad \text{for } |r| \leq s.$$

For every nonnegative integer k , the first occurrence of $-k$ in this sequence is $v_{k^2+1} = -k$, and the first occurrence of k in this sequence is $v_{(k+1)^2} = k$.

The sequence U will be the unique subsequence of V constructed as follows. Let $n \in \mathbf{Z}$. If $f(n) = \infty$, then U will contain the terms $v_{s^2+s+1+n}$ for every $s \geq |n|$. If $f(n) = \ell < \infty$, then U will contain the ℓ terms $v_{s^2+s+1+n}$ for $s = |n|, |n|+1, \dots, |n|+\ell-1$ in the subsequence U , but not the terms $v_{s^2+s+1+n}$ for $s \geq |n|+\ell$. Let $m_1 < m_2 < m_3 < \dots$ be the strictly increasing sequence of positive integers such that $\{v_{m_k}\}_{k=1}^{\infty}$ is the resulting subsequence of V . Let $U = \{u_k\}_{k=1}^{\infty}$, where $u_k = v_{m_k}$. Then

$$f(n) = \text{card}(\{k \geq 1 : u_k = n\}).$$

Let $\text{card}(f^{-1}(0)) = \Delta$. The sequence U also has the following property: If $|u_k| = n$, then for every integer $m \notin f^{-1}(0)$ with $|m| < n$ there is a positive integer $j < k$ with $u_j = m$. It follows that

$$\{0, 1, -1, 2, -2, \dots, n-1, -(n-1)\} \setminus f^{-1}(0) \subseteq \{u_1, u_2, \dots, u_{k-1}\},$$

and so

$$k-1 \geq 2(n-1) + 1 - \Delta.$$

This implies that

$$|u_k| = n \leq \frac{k + \Delta}{2}.$$

Since u_k is an integer, we have

$$|u_k| \leq \left\lceil \frac{k + \Delta}{2} \right\rceil.$$

This completes the proof. $\square$

Lemma 1 is best possible in the sense that for every nonnegative integer Δ there is a function $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ with $\text{card}(f^{-1}(0)) = \Delta$ and a sequence $U = \{u_k\}_{k=1}^{\infty}$ of integers such that

$$|u_k| = \left\lceil \frac{k + \Delta}{2} \right\rceil \quad \text{for all } k \geq 1. \quad (1)$$

For example, if $\Delta = 2\delta + 1$ is odd, define the function f by

$$f(n) = \begin{cases} 0 & \text{if } |n| \leq \delta \\ 1 & \text{if } |n| \geq \delta + 1 \end{cases}$$

and the sequence U by

$$\begin{aligned} u_{2i-1} &= \delta + i, \\ u_{2i} &= -(\delta + i) \end{aligned}$$

for all $i \geq 1$.

If $\Delta = 2\delta$ is even, define f by

$$f(n) = \begin{cases} 0 & \text{if } -\delta \leq n \leq \delta - 1 \\ 1 & \text{if } n \geq \delta \text{ or } n \leq -\delta - 1 \end{cases}$$

and the sequence U by $u_1 = \delta$ and

$$\begin{aligned} u_{2i} &= \delta + i, \\ u_{2i+1} &= -(\delta + i) \end{aligned}$$

for all $i \geq 1$. In both cases the sequence U satisfies (1).

The set A is called a *Sidon set of order h* if $r_{A,h}(n) = 0$ or 1 for every integer n . If A is a Sidon set of order h , then A is a Sidon set of order j for all $j = 1, 2, \dots, h$.

Lemma 2 *Let A be a finite Sidon set of order h and $d = \max(\{|a| : a \in A\})$. If $|c| > (2h - 1)d$, then $A \cup \{c\}$ is also a Sidon set of order h .*

Proof. Let $n \in h(A \cup \{c\})$. Suppose that

$$n = a_1 + \dots + a_j + (h - j)c = a'_1 + \dots + a'_\ell + (h - \ell)c,$$

where

$$\begin{aligned} 0 &\leq j \leq \ell \leq h, \\ a_1, \dots, a_j, a'_1, \dots, a'_\ell &\in A, \end{aligned}$$

and

$$a_1 \leq \dots \leq a_j \quad \text{and} \quad a'_1 \leq \dots \leq a'_\ell.$$

If $j < \ell$, then

$$\begin{aligned} |c| &\leq |(\ell - j)c| \\ &= |a'_1 + \dots + a'_\ell - (a_1 + \dots + a_j)| \\ &\leq (\ell + j)d \\ &\leq (2h - 1)d \\ &< |c|, \end{aligned}$$

which is absurd. Therefore, $j = \ell$ and $a_1 + \dots + a_j = a'_1 + \dots + a'_j$. Since A is a Sidon set of order j , it follows that $a_i = a'_i$ for all $i = 1, \dots, j$. Consequently, $A \cup \{c\}$ is a Sidon set of order h . $\square$

3 Construction of asymptotic bases

We can now construct asymptotic bases of order h for the integers with arbitrary representation functions.

Theorem 1 *Let $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ be a function such that the set $f^{-1}(0)$ is finite. Let $\varphi : \mathbf{N}_0 \rightarrow \mathbf{R}$ be a nonnegative function such that $\lim_{x \rightarrow \infty} \varphi(x) = \infty$. For every $h \geq 2$ there exist infinitely many asymptotic bases A of order h for the integers such that*

$$r_{A,h}(n) = f(n) \quad \text{for all } n \in \mathbf{Z},$$

and

$$A(-x, x) \leq \varphi(x)$$

for all $x \geq 0$.

Proof. By Lemma 1, there is a sequence $U = \{u_k\}_{k=1}^{\infty}$ of integers such that

$$f(n) = \text{card}(\{k \geq 1 : u_k = n\})$$

for every integer n .

Let $h \geq 2$. We shall construct an ascending sequence of finite sets $A_1 \subseteq A_2 \subseteq A_3 \subseteq \dots$ such that, for all positive integers k and for all integers n ,

(i)

$$r_{A_k,h}(n) \leq f(n),$$

(ii)

$$r_{A_k,h}(n) \geq \text{card}(\{i : 1 \leq i \leq k \text{ and } u_i = n\}),$$

(iii)

$$\text{card}(A_k) \leq 2k,$$

(iv)

$$A_k \text{ is a Sidon set of order } h-1.$$

Conditions (i) and (ii) imply that the infinite set

$$A = \bigcup_{k=1}^{\infty} A_k$$

is an asymptotic basis of order h for the integers such that $r_{A,h}(n) = f(n)$ for all $n \in \mathbf{Z}$.

We construct the sets A_k by induction. Since the set $f^{-1}(0)$ is finite, there exists an integer d_0 such that $f(n) \geq 1$ for all integers n with $|n| \geq d_0$. If $u_1 \geq 0$, choose a positive integer $c_1 > 2hd_0$. If $u_1 < 0$, choose a negative integer $c_1 < -2hd_0$. Then

$$|c_1| > 2hd_0.$$

Let

$$A_1 = \{-c_1, (h-1)c_1 + u_1\}.$$

The sumset hA_1 is the finite arithmetic progression

$$\begin{aligned} hA_1 &= \{-hc_1 + (hc_1 + u_1)i : i = 0, 1, \dots, h\} \\ &= \{-hc_1, u_1, hc_1 + 2u_1, 2hc_1 + 3u_1, \dots, (h-1)hc_1 + hu_1\}. \end{aligned}$$

Then $|n| \geq hc_1 > d_0$ for all $n \in hA_1 \setminus \{u_1\}$, and so

$$r_{A_1, h}(n) = 1 \leq f(n)$$

for all $n \in hA_1$. It follows that $r_{A_1, h}(n) \leq f(n)$ for all $n \in \mathbf{Z}$. The set A_1 is a Sidon set of order h , hence also a Sidon set of order $h-1$. Thus, the set A_1 satisfies conditions (i)–(iv).

We assume that for some integer $k \geq 2$ we have constructed a set A_{k-1} satisfying conditions (i) – (iv). If

$$r_{A_{k-1}}(n) \geq \text{card}(\{i : 1 \leq i \leq k \text{ and } u_i = n\})$$

for all $n \in \mathbf{Z}$, then the set $A_k = A_{k-1}$ satisfies conditions (i) – (iv). Otherwise,

$$r_{A_{k-1}}(u_k) = \text{card}(\{i : 1 \leq i \leq k \text{ and } u_i = u_k\}) - 1 < f(u_k).$$

We shall construct a Sidon set A_k of order $h-1$ such that

$$\text{card}(A_k) = \text{card}(A_{k-1}) + 2$$

and

$$r_{A_k, h}(n) = \begin{cases} r_{A_{k-1}, h}(n) + 1 & \text{if } n = u_k \\ r_{A_{k-1}, h}(n) & \text{if } n \in hA_{k-1} \setminus \{u_k\} \\ 1 & \text{if } n \in hA_k \setminus (hA_{k-1} \cup \{u_k\}). \end{cases} \quad (2)$$

Define the integer

$$d_{k-1} = \max(\{|a| : a \in A_{k-1} \cup \{u_k\}\}). \quad (3)$$

Then

$$A_{k-1} \subseteq [-d_{k-1}, d_{k-1}].$$

If $u_k \geq 0$, choose a positive integer c_k such that $c_k > 2hd_{k-1}$. If $u_k < 0$, choose a negative integer c_k such that $c_k < -2hd_{k-1}$. Then

$$|c_k| > 2hd_{k-1}. \quad (4)$$

Let

$$A_k = A_{k-1} \cup \{-c_k, (h-1)c_k + u_k\}.$$

Then

$$\text{card}(A_k) = \text{card}(A_{k-1}) + 2 \leq 2k.$$

We shall assume that $u_k \geq 0$, hence $c_k > 0$. (The argument in the case $u_k < 0$ is similar.) We decompose the sumset hA_k as follows:

$$hA_k = \bigcup_{\substack{r+i+j=h \\ r,i,j \geq 0}} (r(h-1)c_k + ru_k - ic_k + jA_{k-1}) = \bigcup_{r=0}^h B_r,$$

where

$$B_r = r(h-1)c_k + ru_k + \bigcup_{i=0}^{h-r} (-ic_k + (h-r-i)A_{k-1}).$$

If $n \in B_r$, then there exist integers $i \in \{0, 1, \dots, h-r\}$ and $y \in (h-r-i)A_{k-1}$ such that

$$n = r(h-1)c_k + ru_k - ic_k + y.$$

Since

$$|y| \leq (h-r-i)d_{k-1},$$

it follows that

$$n \geq r(h-1)c_k + ru_k - ic_k - (h-r-i)d_{k-1} \quad (5)$$

and

$$n \leq r(h-1)c_k + ru_k - ic_k + (h-r-i)d_{k-1}.$$

Let $m \in B_{r-1}$ and $n \in B_r$ for some $r \in \{1, \dots, h\}$. There exist nonnegative integers $i \leq h-r$ and $j \leq h-r+1$ such that

$$\begin{aligned} n - m &\geq (r(h-1)c_k + ru_k - ic_k - (h-r-i)d_{k-1}) \\ &\quad - ((r-1)(h-1)c_k + (r-1)u_k - jc_k + (h-r+1-j)d_{k-1}) \\ &= (h-1+j-i)c_k + u_k - (2h-2r-i-j+1)d_{k-1} \\ &\geq (h-1-i)c_k - 2hd_{k-1}. \end{aligned}$$

If $r \geq 2$, then $i \leq h-2$ and inequality (4) implies that

$$n - m \geq c_k - 2hd_{k-1} > 0.$$

Therefore, if $m \in B_{r-1}$ and $n \in B_r$ for some $r \in \{2, \dots, h\}$, then $m < n$.

In the case $r = 1$ we have $m \in B_0$ and $n \in B_1$. If $i \leq h-2$, then (4) implies that

$$n - m \geq (h-1-i)c_k - 2hd_{k-1} \geq c_k - 2hd_{k-1} > 0$$

and (5) implies that

$$n \geq (h-1-i)c_k + u_k - (h-1-i)d_{k-1} > c_k - hd_{k-1} > d_0.$$

If $r = 1$ and $i = h-1$, then $n = u_k$. Therefore, if $m \in B_0$ and $n \in B_1$, then $m < n$ unless $m = n = u_k$. It follows that the sets $B_0, B_1 \setminus \{u_k\}, B_2, \dots, B_h$ are pairwise disjoint.

Let $n \in B_r$ for some $r \geq 1$. Suppose that $0 \leq i \leq j \leq h - r$, and that

$$n = r(h - 1)c_k + ru_k - ic_k + y \quad \text{for some } y \in (h - r - i)A_{k-1}$$

and

$$n = r(h - 1)c_k + ru_k - jc_k + z \quad \text{for some } z \in (h - r - j)A_{k-1}.$$

Subtracting these equations, we obtain

$$z - y = (j - i)c_k.$$

Recall that $|a| \leq d_{k-1}$ for all $a \in A_{k-1}$. If $i < j$, then

$$\begin{aligned} c_k &\leq (j - i)c_k = z - y \\ &\leq |y| + |z| \leq (2h - 2r - i - j)d_{k-1} \\ &< 2hd_{k-1} < c_k, \end{aligned}$$

which is impossible. Therefore, $i = j$ and $y = z$. Since $0 \leq h - r - i \leq h - 1$ and A_{k-1} is a Sidon set of order $h - 1$, it follows that

$$r_{A_{k-1}, h-r-i}(y) = 1$$

and so

$$r_{A_k, h}(n) = 1 \leq f(n) \quad \text{for all } n \in (B_1 \setminus \{u_k\}) \cup \bigcup_{r=2}^h B_r.$$

Next we consider the set

$$B_0 = hA_{k-1} \cup \bigcup_{i=1}^h (-ic_k + (h - i)A_{k-1}).$$

For $i = 1, \dots, h$, we have

$$c_k > 2hd_{k-1} \geq (2h - 2i + 1)d_{k-1}$$

and so

$$\begin{aligned} \max(-ic_k + (h - i)A_{k-1}) &\leq -ic_k + (h - i)d_{k-1} \\ &< -(i - 1)c_k - (h - i + 1)d_{k-1} \\ &\leq \min(-(i - 1)c_k + (h - i + 1)A_{k-1}). \end{aligned}$$

Therefore, the sets $-ic_k + (h - i)A_{k-1}$ are pairwise disjoint for $i = 0, 1, \dots, h$. In particular, if $n \in B_0 \setminus hA_{k-1}$, then

$$n \leq \max(-c_k + (h - 1)A_{k-1}) \leq -c_k + (h - 1)d_{k-1} < -d_{k-1} \leq -d_0$$

and $f(n) \geq 1$. Since A_{k-1} is a Sidon set of order $h - 1$, it follows that

$$r_{A_k, h}(n) = 1 \leq f(n) \quad \text{for all } n \in \bigcup_{i=1}^h (-ic_k + (h - i)A_k) = B_0 \setminus hA_{k-1}.$$

It follows from (3) that for any $n \in B_0 \setminus hA_{k-1}$ we have

$$n < -d_{k-1} \leq u_k,$$

and so $u_k \notin B_0 \setminus hA_{k-1}$. Therefore,

$$r_{A_k, h}(u_k) = r_{A_{k-1}, h}(u_k) + 1,$$

and the representation function $r_{A_k, h}$ satisfies the three requirements of (2).

We shall prove that

$$A_k = A_{k-1} \cup \{-c_k, (h-1)c_k + u_k\}.$$

is a Sidon set of order $h-1$. Since A_{k-1} is a Sidon set of order $h-1$ with $d_{k-1} \geq \max\{|a| : a \in A_{k-1}\}$, and since

$$c_k > 2hd_{k-1} > (2(h-1) - 1)d_{k-1},$$

Lemma 2 implies that $A_{k-1} \cup \{-c_k\}$ is a Sidon set of order $h-1$.

Let $n \in (h-1)A_k$. Suppose that

$$\begin{aligned} n &= r(h-1)c_k + ru_k - ic_k + x \\ &= s(h-1)c_k + su_k - jc_k + y, \end{aligned}$$

where

$$\begin{aligned} 0 &\leq r \leq s \leq h-1, \\ 0 &\leq i \leq h-1-r, \\ 0 &\leq j \leq h-1-s, \\ x &\in (h-1-r-i)A_{k-1}, \end{aligned}$$

and

$$y \in (h-1-s-j)A_{k-1}.$$

Then

$$|x| \leq (h-1-r-i)d_{k-1}$$

and

$$|y| \leq (h-1-s-j)d_{k-1}.$$

If $r < s$, then $j \leq h-2$ and

$$\begin{aligned} (h-1)c_k &\leq (s-r)(h-1)c_k + (s-r)u_k \\ &= (j-i)c_k + x - y \\ &\leq (j-i)c_k + (2h-2-r-s-i-j)d_{k-1} \\ &\leq (h-2)c_k + 2hd_{k-1} \\ &< (h-1)c_k, \end{aligned}$$

which is absurd. Therefore, $r = s$ and

$$-ic_k + x = -jc_k + y \in (h - 1 - r)(A_k \cup \{-c_k\}).$$

Since $A_k \cup \{-c_k\}$ is a Sidon set of order $h - 1$, it follows that $i = j$ and that x has a unique representation as the sum of $h - 1 - r - i$ elements of A_k . Thus, A_k is a Sidon set of order $h - 1$.

The set A_k satisfies conditions (i) – (iv). It follows by induction that there exists an infinite increasing sequence $A_1 \subseteq A_2 \subseteq \dots$ of finite sets with these properties, and that $A = \cup_{k=1}^{\infty} A_k$ is an asymptotic basis of order h with representation function $r_{A,h}(n) = f(n)$ for all $n \in \mathbf{Z}$.

Let $A_0 = \emptyset$, and let K be the set of all positive integers k such that $A_k \neq A_{k-1}$. Then

$$A = \cup_{k \in K} A_k = \cup_{k \in K} \{-c_k, (h - 1)c_k\}.$$

For $k \in K$, the only constraints on the choice of the number c_k in the construction of the set A_k were the sign of c_k and the growth condition

$$|c_k| > 2hd_{k-1} \quad \text{for all integers } k \in K.$$

We shall prove that we can construct the asymptotic basis A with counting function $A(-x, x) \leq \varphi(x)$ for all $x \geq 0$. Since $\varphi(x) \rightarrow \infty$ as $x \rightarrow \infty$, for every integer $k \geq 0$ there exists an integer w_k such that

$$\varphi(x) \geq 2k \quad \text{for all } x \geq w_k.$$

We now impose the following additional constraint: Choose c_k such that

$$|c_k| \geq w_k \quad \text{for all integers } k \in K.$$

Then

$$A_1(-x, x) = 0 \leq \varphi(x) \quad \text{for } 0 \leq x < |c_1|$$

and

$$A_1(-x, x) \leq 2 \leq \varphi(x) \quad \text{for } x \geq |c_1| \geq w_1.$$

Suppose that $k \geq 2$ and the set A_{k-1} satisfies $A_{k-1}(-x, x) \leq \varphi(x)$ for all $x \geq 0$. Since

$$A_k \cap (-|c_k|, |c_k|) = A_{k-1} \cap (-|c_k|, |c_k|),$$

it follows that

$$A_k(-x, x) = A_{k-1}(-x, x) \leq \varphi(x) \quad \text{for } 0 \leq x < |c_k|$$

and

$$A_k(-x, x) \leq 2k \leq \varphi(x) \quad \text{for } x \geq |c_k| \geq w_k.$$

This proves by induction that $A_k(-x, x) \leq \varphi(x)$ for all k and x . Since $\lim_{k \rightarrow \infty} |c_k| = \infty$, it follows that for any nonnegative integer x we can choose c_k so that $|c_k| > x$ and

$$A(-x, x) = A_k(-x, x) \leq \varphi(x).$$

For every integer $k \in K$ we had infinitely many choices for the integer c_k to use in the construction of the set A_k , and so there are infinitely many asymptotic bases A with the property that $r_A(n) = f(n)$ for all $n \in \mathbf{Z}$. This completes the proof. $\square$

4 Sums of pairwise distinct integers

Let A be a set of integers and h a positive integer. We define the sumset $h \wedge A$ as the set consisting of all sums of h pairwise distinct elements of A , and the *restricted representation function*

$$\hat{r}_{A,h} : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$$

by

$$\hat{r}_{A,h}(n) = \text{card}(\{\{a_1, \dots, a_h\} \subseteq A : a_1 + \dots + a_h = n \text{ and } a_1 < \dots < a_h\}).$$

The set A of integers is called a *restricted asymptotic basis of order h* if $h \wedge A$ contains all but finitely many integers, or, equivalently, if $\hat{r}_{A,h}^{-1}(0)$ is a finite subset of $\mathbf{Z}$.

The following theorem can also be proved by the method used to prove Theorem 1.

Theorem 2 *Let $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ be a function such that $f^{-1}(0)$ is a finite set of integers. Let $\varphi : \mathbf{N}_0 \rightarrow \mathbf{R}$ be a nonnegative function such that $\lim_{x \rightarrow \infty} \varphi(x) = \infty$. For every $h \geq 2$ there exist infinitely many sets A of integers such that*

$$\hat{r}_{A,h}(n) = f(n) \quad \text{for all } n \in \mathbf{Z}$$

and

$$A(-x, x) \leq \varphi(x)$$

for all $x \geq 0$.

5 Open problems

Let X be an abelian semigroup, written additively, and let A be a subset of X . We define the h -fold sumset hA as the set consisting of all sums of h not necessarily distinct elements of A . The set A is called an *asymptotic basis of order h for X* if the sumset hA consists of all but at most finitely many elements of X . We also define the h -fold *restricted sumset* $h \wedge A$ as the set consisting of all sums of h pairwise distinct elements of A . The set A is called a *restricted asymptotic basis of order h for X* if the restricted sumset $h \wedge A$ consists of all but at most finitely many elements of X . The classical problems of additive number theory concern the semigroups $\mathbf{N}_0$ and $\mathbf{Z}$.

There are four different representation functions that we can associate to every subset A of X and every positive integer h . Let $(a_1, \dots, a_h)$ and $(a'_1, \dots, a'_h)$ be h -tuples of elements of X . We call these h -tuples *equivalent* if there is a permutation σ of the set $\{1, \dots, h\}$ such that $a'_{\sigma(i)} = a_i$ for all $i = 1, \dots, h$. For every $x \in X$, let $r_{A,h}(x)$ denote the number of equivalence classes of h -tuples $(a_1, \dots, a_h)$ of elements of A such that $a_1 + \dots + a_h = x$. The function $r_{A,h}$ is called the *unordered representation function* of A . This is the function that we studied in this paper. The set A is an asymptotic basis of order h if $r_{A,h}^{-1}(0)$ is a finite subset of X .

Let $R_{A,h}(x)$ denote the number of h -tuples $(a_1, \dots, a_h)$ of elements of A such that $a_1 + \dots + a_h = x$. The function $R_{A,h}$ is called the *ordered representation function* of A .

Let $\hat{r}_{A,h}(x)$ denote the number of equivalence classes of h -tuples $(a_1, \dots, a_h)$ of pairwise distinct elements of A such that $a_1 + \dots + a_h = x$, and let $\hat{R}_{A,h}(x)$ denote the number of h -tuples $(a_1, \dots, a_h)$ of pairwise distinct elements of A such that $a_1 + \dots + a_h = x$. These functions are called the *unordered restricted representation function* of A and the *ordered restricted representation function* of A , respectively. The two restricted representation functions are essentially identical, since $\hat{R}_{A,h}(x) = h! \hat{r}_{A,h}(x)$ for all $x \in X$.

In the discussion below, we use only the unordered representation function $r_{A,h}$, but each of the problems can be reformulated in terms of the other representation functions.

For every countable abelian semigroup X , let $\mathcal{F}(X)$ denote the set of all functions $f : X \rightarrow \mathbf{N}_0 \cup \{\infty\}$, and let $\mathcal{F}_0(X)$ denote the set of all functions $f : X \rightarrow \mathbf{N}_0 \cup \{\infty\}$ such that $f^{-1}(0)$ is a finite subset of X . Let $\mathcal{F}_c(X)$ denote the set of all functions $f : X \rightarrow \mathbf{N}_0 \cup \{\infty\}$ such that $f^{-1}(0)$ is a cofinite subset of X , that is, $f(x) \neq 0$ for only finitely many $x \in X$, or, equivalently,

$$\text{card}(f^{-1}(\mathbf{N} \cup \{\infty\})) < \infty.$$

Let $\mathcal{R}(X, h)$ denote the set of all h -fold representation functions of subsets A of X . If $r_{A,h}$ is the representation function of an asymptotic basis A of order h for X , then $r_{A,h}^{-1}(0)$ is a finite subset of X , and so $r_{A,h} \in \mathcal{F}_0(X)$. Let $\mathcal{R}_0(X, h)$ denote the set of all h -fold representation functions of asymptotic bases A of order h for X . Let $\mathcal{R}_c(X, h)$ denote the set of all h -fold representation functions of finite subsets of X . We have

$$\mathcal{R}(X, h) \subseteq \mathcal{F}(X),$$

$$\mathcal{R}_0(X, h) \subseteq \mathcal{F}_0(X),$$

and

$$\mathcal{R}_c(X, h) \subseteq \mathcal{F}_c(X),$$

In the case $h = 1$, we have, for every set $A \subseteq X$,

$$r_{A,1}(x) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{if } x \notin A, \end{cases}$$

and so

$$\begin{aligned}\mathcal{R}(X, 1) &= \{f : X \rightarrow \{0, 1\}\}, \\ \mathcal{R}_0(X, 1) &= \{f : X \rightarrow \{0, 1\} : \text{card}(f^{-1}(0)) < \infty\},\end{aligned}$$

and

$$\mathcal{R}_c(X, 1) = \{f : X \rightarrow \{0, 1\} : \text{card}(f^{-1}(\mathbf{N} \cup \{\infty\})) < \infty\}.$$

In this paper we proved that

$$\mathcal{R}_0(\mathbf{Z}, h) = \mathcal{F}_0(\mathbf{Z}) \quad \text{for all } h \geq 2.$$

Nathanson [8] has extended this result to certain countably infinite groups and semigroups. Let G be any countably infinite abelian group such that $\{2g : g \in G\}$ is infinite. For the unordered restricted representation function $\hat{r}_{A,2}$, we have

$$\mathcal{R}_0(G, 2) = \mathcal{F}_0(G).$$

More generally, let S is any countable abelian semigroup such that for every $s \in S$ there exist $s', s'' \in S$ with $s = s' + s''$. In the abelian semigroup $X = S \oplus G$, we have

$$\mathcal{R}_0(X, 2) = \mathcal{F}_0(X).$$

If $\{12g : g \in G\}$ is infinite, then $\mathcal{R}_0(X, 2) = \mathcal{F}_0(X)$ for the unordered representation function $r_{A,2}$.

The following problems are open for all $h \geq 2$:

1. Determine $\mathcal{R}_0(\mathbf{N}_0, h)$. Equivalently, describe the representation functions of additive bases for the nonnegative integers. This is a major unsolved problem in additive number theory, of which the Erdős-Turán conjecture is only a special case.
2. Determine $\mathcal{R}(\mathbf{Z}, h)$. In this paper we computed $\mathcal{R}_0(\mathbf{Z}, h)$, the set of representation functions of additive bases for the integers, but it is not known under what conditions a function $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ with $f^{-1}(0)$ infinite is the representation function of a subset A of X . It can be proved that if $f^{-1}(0)$ is infinite but sufficiently sparse, then $f \in \mathcal{R}(\mathbf{Z}, h)$.
3. Determine $\mathcal{R}(\mathbf{N}_0, h)$. Is there a simple list of necessary and sufficient conditions for a function $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ to be the representation function of some set of nonnegative integers?
4. Determine $\mathcal{R}_c(\mathbf{Z}, h)$. Describe the representation functions of finite sets of integers. If A is a finite set of integers and t is an integer, then for the translated set $t + A$ we have

$$r_{t+A,h}(n) = r_{A,h}(n - ht)$$

for all integers n . This implies that if $f(n) \in \mathcal{R}_c(\mathbf{Z}, h)$, then $f(n - ht) \in \mathcal{R}_c(\mathbf{Z}, h)$ for every integer b , so it suffices to consider only finite sets A of nonnegative integers with $0 \in A$, and functions $f \in \mathcal{F}_c(\mathbf{N}_0, h)$ with $f(0) = 1$.

5. Determine $\mathcal{R}_0(G, 2)$, $\mathcal{R}(G, 2)$, and $\mathcal{R}_c(G, 2)$ for the infinite abelian group $G = \oplus_{i=1}^{\infty} \mathbf{Z}/2\mathbf{Z}$. Note that $\{2g : g \in G\} = \{0\}$ for this group.
6. Determine $\mathcal{R}_0(G, h)$ and $\mathcal{R}(G, h)$, where G is an arbitrary countably infinite abelian group and $h \geq 2$.
7. There is a class of problems of the following type. Do there exist integers h and k with $2 \leq h < k$ such that

$$\mathcal{R}(\mathbf{Z}, h) \neq \mathcal{R}(\mathbf{Z}, k)?$$

We can easily find sets of integers to show that that $\mathcal{R}_0(\mathbf{N}_0, h) \neq \mathcal{R}_0(\mathbf{N}_0, k)$. For example, let $A = \mathbf{N}$ be the set of all positive integers, and let $h \geq 1$. Then $r_{\mathbf{N},h}(0) = 0$ and $r_{\mathbf{N},h}(h) = 1$. If B is any set of nonnegative integers and $k > h$, then either $r_{B,k}(0) = 1$ or $r_{B,k}(h) = 0$, and so $r_{\mathbf{N},h} \notin \mathcal{R}_0(\mathbf{N}_0, k)$. Is it true that

$$\mathcal{R}_0(\mathbf{N}_0, h) \cap \mathcal{R}_0(\mathbf{N}_0, k) = \emptyset$$

for all $h \neq k$?

8. By Theorem 1, for every $h \geq 2$ and every function $f \in \mathcal{F}_0(\mathbf{Z})$, there exist arbitrarily sparse sets A of integers such that $r_{A,h}(n) = f(n)$ for all n . It is an open problem to determine how dense the sets A can be. For example, in the special case $h = 2$ and $f(n) = 1$, Nathanson [7] proved that there exists a set A such that $r_{A,2}(n) = 1$ for all n , and $\log x \ll A(-x, x) \ll \log x$. For an arbitrary representation function $f \in \mathcal{F}_0(\mathbf{Z})$, Nathanson [6] constructed an asymptotic basis of order h with $A(-x, x) \gg x^{1/(2h-1)}$. In the case $h = 2$, Cilleruelo and Nathanson [1] improved this to $A(-x, x) \gg x^{\sqrt{2}-1}$.

References

- [1] J. Cilleruelo and M. B. Nathanson, *Dense sets of integers with prescribed representation functions*, in preparation.
- [2] G. A. Dirac, *Note on a problem in additive number theory*, J. London Math. Soc. **26** (1951), 312–313.
- [3] P. Erdős and P. Turán, *On a problem of Sidon in additive number theory and some related questions*, J. London Math. Soc. **16** (1941), 212–215.
- [4] P. Erdős and W. H. J. Fuchs, *On a problem of additive number theory*, J. London Math. Soc. **31** (1956), 67–73.
- [5] M. B. Nathanson, *Representation functions of sequences in additive number theory*, Proc. Amer. Math. Soc. **72** (1978), 16–20.

- [6] ———, *The inverse problem for representation functions of additive bases*, Number Theory: New York 2003 (New York), Springer-Verlag, 2004.
- [7] ———, *Unique representation bases for the integers*, Acta Arith. **108** (2003), no. 1, 1–8.
- [8] ———, *Representation functions of additive bases for abelian semigroups*, Int. J. Math. Math. Sci. (2004), to appear.